

Assistant Professor of Orthopaedics Division of Sports Medicine Tel: (646) 501-7223

## **Distal Biceps Tendon Repair Rehabilitation Guidelines**

Name:	Date:
Diagnosis:	Date of Surgery:
<ul> <li>IMMOBILIZATION</li> <li>Posterior splint</li> <li>Elbow immobilization at 90 degrees for 7-10 day</li> </ul>	s
<ul> <li>BRACE</li> <li>Elbow placed in ROM brace at 7-10 days post-ope</li> <li>ROM set at 45 degrees to full flexion</li> <li>Gradually increase elbow ROM in brace</li> </ul>	erative
<ul> <li>RANGE OF MOTION PROGRESSION</li> <li>Week 2: 45 degrees to full flexion</li> <li>Week 3: 45 degrees to full flexion</li> <li>Week 4: 30 degrees to full flexion</li> <li>Week 5: 20 degrees to full flexion</li> <li>Week 6: 10 degrees to full flexion; full supination</li> <li>Week 8: full ROM of elbow; full supination/prona</li> </ul>	• •
<ul> <li>RANGE OF MOTION EXERCISES</li> <li>Week 3-5: passive ROM for elbow flexion and suppronation</li> <li>Week 6-8: initiate AAROM elbow flexion</li> <li>Week 9: active ROM elbow flexion</li> </ul>	oination; active-assisted ROM for elbow extension and
<ul> <li>STRENGTHENING PROGRAM</li> <li>Week 2-3: isometrics for triceps and shoulder mu</li> <li>Week 9: active ROM, no resistance applied</li> <li>Week 10: PRE (progressive resistive exercise) prosupination/pronation         <ul> <li>Begin with one pound and gradually increase.</li> </ul> </li> <li>Week 14: may initiate light weight training such an exercise.</li> </ul>	ogram is initiated for elbow flexion and ease shoulder strengthening program
Comments:	
Frequency: times per week Duration:	weeks
Signature:	Date: